



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2018

---

## **Agrobacterium-mediated floral dip transformation of the model polyploid species *Arabidopsis kamchatica***

Yew, Chow-Lih ; Kakui, Hiroyuki ; Shimizu, Kentaro K

**Abstract:** Polyploidization has played an important role in the speciation and diversification of plant species. However, genetic analyses of polyploids are challenging because the vast majority of the model species are diploids. The allotetraploid *Arabidopsis kamchatica*, which originated through the hybridization of the diploid *Arabidopsis halleri* and *Arabidopsis lyrata*, is an emerging model system for studying various aspects of polyploidy. However, a transgenic method that allows the insertion of a gene of interest into *A. kamchatica* is still lacking. In this study, we investigated the early development of pistils in *A. kamchatica* and confirmed the formation of open pistils in young flower buds (stages 8–9), which is important for allowing *Agrobacterium* to access female reproductive tissues. We established a simple *Agrobacterium*-mediated floral dip transformation method to transform a gene of interest into *A. kamchatica* by dipping *A. kamchatica* inflorescences bearing many young flower buds into a 5% sucrose solution containing 0.05% Silwet L-77 and *Agrobacterium* harboring the gene of interest. We showed that a screenable marker comprising fluorescence-accumulating seed technology with green fluorescent protein was useful for screening the transgenic seeds of two accessions of *A. kamchatica* subsp. *kamchatica* and an accession of *A. kamchatica* subsp. *kawasakiana*.

DOI: <https://doi.org/10.1007/s10265-017-0982-9>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-141199>

Journal Article

Accepted Version

Originally published at:

Yew, Chow-Lih; Kakui, Hiroyuki; Shimizu, Kentaro K (2018). *Agrobacterium*-mediated floral dip transformation of the model polyploid species *Arabidopsis kamchatica*. *Journal of Plant Research*, 131(2):349–358.

DOI: <https://doi.org/10.1007/s10265-017-0982-9>

**Electric supplementary materials**

**Title:**

***Agrobacterium*-mediated floral dip transformation of the model polyploid species**

***Arabidopsis kamchatica***

**Authors:**

Chow-Lih Yew<sup>1</sup>, Hiroyuki Kakui<sup>1,2</sup> & Kentaro K. Shimizu<sup>1,2</sup>

<sup>1</sup>Department of Evolutionary Biology and Environmental Studies, University of Zurich, Winterthurerstrasse 190, 8057 Zurich, Switzerland.

<sup>2</sup> Kihara Institute for Biological Research, Yokohama City University, 641-12 Maioka, 244-0813 Totsuka-ward, Yokohama, Japan.

**Journal:**

Journal of Plant Research

**Corresponding author:**

Kentaro K. Shimizu

Mailing address: Department of Evolutionary Biology and Environmental Studies,  
University of Zurich, Winterthurerstrasse 190, 8057 Zurich, Switzerland.


Phone number: +41 44 635 6740

E-mail address: kentaro.shimizu@ieu.uzh.ch


**Content:**

**Figs. S1–S4**


**Table S1**




**Fig. S1** Primers designed in the pFAST-G01-geneA binary vector for PCR amplification. Black arrows, specific primers designed for PCR amplification (Table S1); *OLE1*, *OLEOSIN1*; *pOLE1*, promoter of *OLE1*; *GFP*, green fluorescent protein; *Hyg*, hygromycin resistance gene; *Sm/SpR*, spectinomycin resistance gene; LB, left border; RB, right border; t35S, terminator of CaMV 35S; attB1 and attB2, Gateway recombination sequences.



**Fig. S2** Observation of seeds from the Takashima accession under a fluorescence stereomicroscope. **a** Takashima accession seeds exhibited strong autofluorescence under a fluorescence stereomicroscope equipped with an RFP filter. **b** Takashima accession seeds exhibited weak autofluorescence under a fluorescence stereomicroscope equipped with a GFP filter. Bar, 1 mm.



**Fig. S3** Kanamycin treatment of the wild-type Potter accessions and T<sub>1</sub> seeds of transgenic *A. thaliana* transformed with the kanamycin resistance gene as a control. Seeds were grown on ½× MS plates containing: **a, b** 50 µg ml<sup>-1</sup> and **c, d** 250 µg ml<sup>-1</sup> of kanamycin. Bar, 2 cm.



**Fig. S4** Treatment with Basta for 2–3 weeks killed the Takashima accession seedlings.

**Table S1** Specific primers designed in the pFAST-G01-geneA binary vector for PCR amplification

Forward primer	Sequence	Reverse primer	Sequence	Annealing temperature (°C)	Elongation time (s)
GFP_F	ACGTAAACGG CCACAAGTTC	GFP_R	TGTGGCGGATC TTGAAGTTCAC	60	30
Hyg_F	GGGGAATTTAT GGAACGTCA	pOLE1_R	CTCTAGAGGAT CCCCGGGTA	60	30